Draft version October 12, 2010
Preprint typeset using LaTeX style emulateapj v. 08/22/09

arXiv:1010.2093v1 [astro-ph.GA] 11 Oct 2010

PHOTOCHEMISTRY IN THE INNER LAYERS OF CLUMPY CIRCUMSTELLAR ENVELOPES: FORMATION OF WATER IN C-RICH OBJECTS AND OF C-BEARING MOLECULES IN O-RICH OBJECTS

Marcelino Agúndez[1], José Cernicharo[2] and Michel Guélin[3]

Draft version October 12, 2010

ABSTRACT

A mechanism based on the penetration of interstellar ultraviolet photons into the inner layers of clumpy circumstellar envelopes around AGB stars is proposed to explain the non-equilibrium chemistry observed in such objects. We show through a simple modelling approach that in circumstellar envelopes with a certain degree of clumpiness or with moderately low mass loss rates (a few 10^{-7} $M_\odot$ yr^{-1}) a photochemistry can take place in the warm and dense inner layers inducing important changes in the chemical composition. In carbon-rich objects water vapor and ammonia would be formed with abundances of 10^{-8}–10^{-6} relative to H_2, while in oxygen-rich envelopes ammonia and carbon-bearing molecules such as HCN and CS would form with abundances of 10^{-9}–10^{-7} relative to H_2. The proposed mechanism would explain the recent observation of warm water vapor in the carbon-rich envelope IRC +10216 with the Herschel Space Observatory, and predict that H_2O should be detectable in other carbon-rich objects.

Subject headings: astrochemistry — molecular processes — stars: AGB and post-AGB — circumstellar matter

1. INTRODUCTION

Evolved stars in the asymptotic giant branch (AGB) phase undergo important mass loss processes that produce extended circumstellar envelopes (CSEs) composed of dust and molecules. The molecular composition of CSEs is affected by several processes during the travel towards the interstellar medium (ISM) but it is originally established in the stellar atmosphere under thermochemical equilibrium (TE) conditions and is to a large extent governed by the C/O abundance ratio. In C-rich stars (C/O > 1) most of the oxygen is in the form of CO resulting in a carbon-based chemistry while in O-rich stars (C/O < 1) CO locks most of the carbon leaving little for other molecules (Tsuji 1973; see also Fig. 1).

Astronomical observations have in the main confirmed this picture, although with a significant number of discrepancies. HCN emission is widely observed in O-rich objects (Bujarrabal et al. 1994; Bieging et al. 2000; Decin et al. 2008), in some of them coming from the inner regions of the CSE (Duari & Hatchell 2000). Water vapor in the C-rich envelope IRC +10216 is confined to the warm near surroundings of the star, according to recent Herschel observations (Decin et al. 2010). Moreover, ammonia is observed in the inner regions of C- and O-rich CSEs (Keady & Ridgway 1993; Menten & Alcolea 1995) with abundances much larger than predicted by TE.

On the theoretical side, shocks induced by the stellar pulsation have been proposed to explain the non-equilibrium chemistry observed in the inner regions of CSEs (Cherchneff 2006). These models explain the formation of HCN and CS in the inner wind of O-rich CSEs, although fail to explain the presence of H_2O in IRC +10216 and of NH_3 in both C- and O-rich CSEs.

In this Letter we investigate an alternative mechanism of non-equilibrium chemistry, based on the penetration of interstellar ultraviolet (UV) photons into the inner regions of CSEs with a certain degree of clumpiness.

2. MODEL

The model is based on a central AGB star surrounded by a spherical circumstellar envelope, whose physical parameters are given in Table 1, and is adopted to investigate the chemistry in both C- and O-rich CSEs with mass loss rates between 10^{-7} and 10^{-5} $M_\odot$ yr^{-1}.

In CSEs with an intense mass loss process strictly isotropic and homogeneous, the inner regions are well shielded from interstellar UV light, so that only the outer layers are affected by photochemistry (e.g. Charnley et al. 1995; Willacy & Millar 1997). Observations, however, have shown that CSEs have usually clumpy structures both at small and large scales (Chapman et al. 1994; Guélin et al. 1997; Weigelt et al. 1998; Fong et al. 2003; Leão et al. 2006), which allow for a deeper penetration of interstellar UV photons into the inner regions. To model the effects of clumpiness on the circumstellar chemistry we adopt a simple approach in which the CSE consists of two components: a major one whose inner regions are well shielded against interstellar UV light, and a minor one (which accounts for a fraction f_m of the total circumstellar mass) for which the shielding matter located in the radial outward direction is grouped into clumpy structures leaving a fraction f_Ω of the solid angle of arrival of interstellar photons free of matter.

The gas phase chemical composition of these two components is computed as they expand from the innermost regions ($r = 2R_*$) until the end of the CSE. The adopted abundances at the initial radius are given in Table 2. ^{13}CO is also included with an abundance 30 times lower than ^{12}CO (Milam et al. 2009). The chemical network has been used in previous chemical models

---

[1] LUTH, Observatoire de Paris-Meudon, 5 Place Jules Janssen, 92190 Meudon, France; marcelino.agundez@obspm.fr
[2] Departamento de Astrofísica, Centro de Astrobiología, CSIC-INTA, Ctra. de Torrejón a Ajalvir km 4, Torrejón de Ardoz, 28850 Madrid, Spain
[3] Institut de Radioastronomie Millimétrique, 300 rue de la Piscine, 38406 Saint Martin d'Héres, France

TABLE 1
Model physical parameters

Star effective temperature (T_*)	2000 K
Stellar radius (R_*)	5×10^{13} cm
Expansion velocity (v)^a	5 km s^{-1} for $r < 5\ R_*$ 15 km s^{-1} for $r \geq 5\ R_*$
Gas kinetic temperature (T)^b	$T_*(r/R_*)^{-0.7}$
Volume density of gas particles (n)^c	$\dot{M}/(4\pi r^2 \langle m_g \rangle v)$

Note. — r is the radius measured from the center of the star. [a] Velocity field similar to that adopted in previous studies of inner CSEs (Keady & Ridgway 1993; Fonfría et al. 2008). [b] Values of the exponent are typically between −0.5 and −1 (Justtanont et al. 1994; Fonfría et al. 2008). [c] $\dot{M}$ is the mass loss rate and $\langle m_g \rangle$ is the mean mass of the gas particles (typically ∼ 2.3 times the mass of the hydrogen atom in CSEs).

TABLE 2
Initial abundances relative to H_2 in C- and O-rich CSEs

Carbon-rich			Oxygen-rich		
Species	Abundance	Ref	Species	Abundance	Ref
He	0.17		He	0.17	
CO	8×10^{-4}	(1)	CO	3×10^{-4}	(1)
N_2	4×10^{-5}	(2)	N_2	4×10^{-5}	(2)
C_2H_2	8×10^{-5}	(3)	H_2O	3×10^{-4}	(4)
HCN	2×10^{-5}	(3)	CO_2	3×10^{-7}	(5)
SiO	1.2×10^{-7}	(3)	SiO	1.7×10^{-7}	(6)
SiS	10^{-6}	(3)	SiS	2.7×10^{-7}	(7)
CS	5×10^{-7}	(3)	SO	10^{-6}	(8)
SiC_2	5×10^{-8}	(3)	H_2S	7×10^{-8}	(9)
HCP	2.5×10^{-8}	(3)	PO	9×10^{-8}	(10)

References. — (1) Teyssier et al. (2006); (2) TE abundance; (3) abundance in IRC +10216, see Agúndez (2009); (4) Maercker et al. (2008); (5) Tsuji et al. (1997); (6) Schöier et al. (2006); (7) Schöier et al. (2007); (8) Bujarrabal et al. (1994); (9) Ziurys et al. (2007); (10) Tenenbaum et al. (2007).

of warm and dense UV illuminated regions (Cernicharo 2004; Agúndez et al. 2008a). Photodissociation and photoionization rates are evaluated as a function of the visual extinction A_V (Woodall et al. 2007; van Dishoeck et al. 2006), adopting the interstellar UV field of Draine (1978). For ^{12}CO the photodissociation rate is evaluated according to Mamon et al. (1988), who included the effect of self-shielding, and for ^{13}CO through the expression $2 \times 10^{-10} \exp(-2.5 A_V)$ s^{-1} (Woodall et al. 2007).

For the major component, shielded by a smooth envelope, A_V depends on the column density of H nuclei $N_{\rm H}$ in the radial outward direction as $A_V = N_{\rm H}({\rm cm}^{-2})/1.87 \times 10^{21}$ (Bohlin et al. 1978). The UV field for this component may be expressed as:

$$4\pi J_\lambda = I_\lambda \Omega \exp\left\{ - \frac{[A_\lambda/A_V]}{1.086} A_V \right\} \qquad (1)$$

where I_λ is the unattenuated interstellar UV field at a wavelength λ, Ω the solid angle of arrival of most of the UV flux (depends strongly on the radius and ranges from a small fraction of π sr in the inner regions up to almost 4π sr in the outermost layers), and $[A_\lambda/A_V]$ the ratio of the dust extinction at λ and at visual wavelengths (3.8 for λ = 1250 Åas found for the ISM by Fitzpatrick & Massa 1990). For the minor component, shielded by a clumpy envelope, the UV field may be analogously expressed as:

$$4\pi J_\lambda = I_\lambda \Omega \Big[(1 - f_\Omega) \exp\left\{ - \frac{[A_\lambda/A_V]}{1.086} A_V \right\} + f_\Omega \Big] \qquad (2)$$

For the minor component we compute an effective visual extinction $A_V^{\rm ef}$ by substituting A_V by $A_V^{\rm ef}$ into Eq. (1) and equalling the right parts of Eqs. (1) and (2).

Near-infrared interferometry, able to trace the dust emission at milli-arcsecond scales, has revealed a extremely clumpy structure in objects such as IRC +10216, with five individual clumps in the close surroundings of the star, some of them with angle separations of up to 20−30° (Weigelt et al. 1998). Millimeter-wave interferometry of molecular lines can, unlike infrared observations of dust, provide information on the projected velocity in the line of sight and allow to build three dimensional maps. Observations of IRC +10216 in molecular lines of CN, C_2H, C_4H, and HC_5N (Guélin et al. 1993; Dinh-V-Trung & Lim 2008) have shown that these species are distributed in a circumstellar shell with a radius of 15-20″, and with two conical holes in the NNE and SSW directions having an aperture angle of about 30°. This corresponds to a solid angle of $\pi/4$ sr, which may be a good fraction of Ω in the inner circumstellar regions. Both f_m and f_Ω are phenomenological parameters, difficult to quantify in any CSE. Anyway, adopting $f_m = 0.1{-}0.2$ (the minor UV illuminated component accounting for just a 10−20 % of the total circumstellar mass) and $f_\Omega = 0.2{-}0.5$ (in line with the observational data described above for IRC +10216) seems reasonable for a CSE with a sufficient degree of clumpiness and allows for a sufficient penetration of interstellar UV photons into the inner layers of the CSE. For all the models we have adopted $f_m = 0.1$ and $f_\Omega = 0.25$.

3. RESULTS

Fig. 1 shows the TE abundance of various molecules calculated under conditions typical of the stellar atmosphere of AGB stars. The calculations have been done with the code described in Tejero & Cernicharo (1991) and most of the thermochemical data have been



Fig. 1.— TE abundances as a function of temperature for a C- and O-rich gas (with a C/O abundance ratio of 1.5 and 0.5 respectively) with a constant particle density of 10^{14} cm^{-3}.



FIG. 2.— Calculated abundance of several molecules as a function of radius for carbon-rich CSEs with mass loss rates of 10^{-5}, 10^{-6}, and 10^{-7} M$_\odot$ yr^{-1}. Dashed-dotted lines correspond to the abundance of the minor UV exposed component properly corrected by the factor f_m, while continuous lines correspond to the abundance weighted-averaged over the minor and major components, which can be expressed as $\overline{x}_i(r) = (1 - f_m) x_i^M(r) + f_m\, x_i^m(r)$, where $x_i^M(r)$ and $x_i^m(r)$ are the abundances of the species i at a radius r in the major and minor components, respectively. Although in the reality the minor UV exposed and the major UV shielded components would not coexist in space, this treatment allows to simplify taking the abundance averaged over all radial directions. Note that in those regions where the dashed-dotted and continuous lines of a given species coincide, only the minor UV exposed component contributes to the abundance.

taken from Chase (1998). The main purpose of these calculations is to show that several molecules which are observed in the inner regions of CSEs have very low TE abundances (e.g. H_2O, NH_3, and SiH_4 in C-rich objects, and HCN, CS, and NH_3 in O-rich objects).

Now focusing on the models based on chemical kinetics, Fig. 2 and Fig. 3 show the calculated abundance distribution of some molecules in C- and O-rich CSEs with mass loss rates of 10^{-5}, 10^{-6}, and 10^{-7} M$_\odot$ yr^{-1}.

In C-rich CSEs with mass loss rates as high as 10^{-5} M$_\odot$ yr^{-1}, typical of an object such as IRC +10216, water vapor would be effectively formed in the dense and warm inner regions of the minor UV illuminated component, for which $A_V^{\rm ef}$ is < 1 mag, with a global abundance relative to H_2 in excess of 10^{-7} (see Fig. 2). In such regions the photodissociation of ^{13}CO and SiO, the major reservoirs of oxygen besides ^{12}CO (hard to photodissociate due to self-shielding effects), liberates atomic oxygen which is effectively converted into water through the chemical reactions

$$O + H_2 \rightarrow OH + H \tag{3}$$

$$OH + H_2 \rightarrow H_2O + H \tag{4}$$

which despite having activation barriers are rapid enough due to the high temperatures attained in these inner layers. The photodestruction of molecules in the minor UV illuminated component occurs fast, but for some of them (e.g. H_2O) the formation rate is high enough to allow them to extend up to relatively large radii, $\sim 10^{16}$ cm (see Fig.2). This mechanism would explain the observation with Herschel of warm water vapor in the inner circumstellar regions of the carbon star IRC +10216 (Decin et al. 2010). Other mechanisms proposed such as sublimation of cometary ices (Melnick et al. 2001), Fischer-Tropsch catalysis on the surface on iron grains (Willacy 2004), or radiative association between O and H_2 (Agúndez & Cernicharo 2006), place water in cool regions located farther than 10^{15} cm.

The proposed mechanism would also have some other interesting consequences. Hydrides other than H_2O, such as NH_3, CH_4, H_2S, SiH_4, and PH_3, could also be effectively formed in the inner CSE by successive hydrogenation reactions of the heavy atom. All them are observed in IRC +10216 (Keady & Ridgway 1993; Cernicharo et al. 2000; Agúndez et al. 2008b) with abundances which, except for CH_4, are much larger than predicted by TE (see Fig. 1). Ammonia, for example, is observed in the inner CSE of IRC +10216 with an abundance relative to H_2 of 10^{-7}–10^{-6} (Keady & Ridgway 1993; Hasegawa et al. 2006), and yet no efficient formation mechanism has been proposed, apart from the suggestion that it could be formed on grain surfaces (Keady & Ridgway 1993). Our model predicts an effective formation for NH_3 and H_2S (see Fig. 2), but not for SiH_4 and PH_3 (likely due to the lack of chemical kinetics data for the relevant hydrogenation reactions). Other molecules such as HC_3N increase their abundance in the inner envelope due to the penetration of interstellar UV photons (see Fig. 2), something that has been recently confirmed through observations of IRC +10216 at $\lambda = 0.9$ mm with the IRAM 30-m telescope (Decin et al. 2010; Kahane et al. in preparation).

Still focusing on C-rich sources (see Fig. 2), if we move toward lower mass loss rates then the whole CSE starts to be more transparent to interstellar UV photons. For example, for a mass loss rate as low as 10^{-7} M$_\odot$ yr^{-1} most of H_2O is formed in the major UV shielded component, which is no longer shielded as it has a visual extinction < 1 mag. Thus, for C-rich CSEs with moderately low mass loss rates (up to a few 10^{-7} M$_\odot$ yr^{-1}) we should expect a relatively large H_2O abundance even if the CSE is not particularly clumpy, prediction that should be easily tested with Herschel.

In the case of O-rich CSEs the penetration of interstellar UV photons into the inner layers has also interesting chemical effects (see Fig. 3). Among them it is worth



FIG. 3.— Same as in Fig. 2 but for oxygen-rich CSEs.

mentioning the formation of NH_3, CH_4, HCN, and CS in the inner envelope with abundances relative to H_2 in the range 10^{-8}–10^{-7}, i.e. much larger than predicted by TE calculations (see Fig. 1). The formation of C-bearing molecules in a dense and warm UV illuminated O-rich gas has been discussed by Agúndez et al. (2008a) in the context of the chemistry of protoplanetary disks. HCN, CS, and NH_3 are observed in O-rich CSEs with abundances of 10^{-7}–10^{-6} relative to H_2 (Bujarrabal et al. 1994; Menten & Alcolea 1995; Bieging et al. 2000), which are somewhat higher than predicted by us. Other mechanisms based on shocks induced by the stellar pulsation (Duari et al. 1999; Cherchneff 2006) predict fractional abundances for HCN and CS of 10^{-6}–10^{-5}, which are somewhat higher than observed, but a negligible abundance for NH_3.

4. CONCLUSIONS

We have shown through a simple modelling approach that in CSEs envelopes with a certain degree of clumpiness or with moderately low mass loss rates (a few 10^{-7} M$_\odot$ yr^{-1}) a photochemistry can take place in the warm and dense inner layers inducing important changes in the chemical composition. This mechanism allows for the formation of H_2O and NH_3 in C-rich objects and HCN, CS, and NH_3 in O-rich objects, with abundances much higher than predicted by thermochemical equilibrium but close to the values typically derived from astronomical observations. This mechanism explains the recent observation of warm water vapor in the carbon-rich envelope IRC +10216 with the Herschel Space Observatory, and predict that H_2O should be detectable in other carbon-rich objects.

M.A. is supported by a *Marie Curie Intra-European Individual Fellowship* within the European Community 7th Framework Programme under grant agreement n° 235753.

REFERENCES

Agúndez, M. & Cernicharo, J. 2006, ApJ, 650, 374
Agúndez, M., Cernicharo, J., & Goicoechea, J. R. 2008a, A&A, 483, 831
Agúndez, M., et al. 2008b, A&A, 485, L33
Agúndez, M. 2009, PhD Thesis, Univ. Autónoma de Madrid
Bieging, J. H., Shaked, S., & Gensheimer, P. D. 2000, ApJ, 543, 897
Bohlin, R. C., Savage, B. D., & Drake, J. F. 1978, ApJ, 224, 132
Bujarrabal, V., Fuente, A., & Omont, A. 1994, A&A, 285, 247
Cernicharo, J., Guélin, M., & Kahane, C. 2000, A&AS, 142, 181
Cernicharo, J. 2004, ApJ, 608, L41
Chapman, J. M., Sivagnanam, P., Cohen, R. J., & Le Squeren, A. M. 1994, MNRAS, 268, 475
Charnley, S. B., Tielens, A. G. G. M., & Kress, M. E. 1995, MNRAS, 274, L53
Chase, M., W. 1998, NIST-JANAF thermochemical Tables, 4th ed., J. Phys. Chem. Ref. Data, Monograph n 9
Cherchneff, I. 2006, A&A, 456, 1001
Decin, L., et al. 2008, A&A, 480, 431
Decin, L., et al. 2010, Nature, 467, 64
Dinh-V-Trung & Lim, J. 2008, ApJ, 678, 303
Draine, B. T. 1978, ApJS, 36, 595
Duari, D., Cherchneff, I., & Willacy, K. 1999, A&A, 341, L47
Duari, D. & Hatchell, J. 2000, A&A, 358, L25
Fitzpatrick, E. L. & Massa, D. 1990, ApJS, 72, 163
Fonfría, J. P., Cernicharo, J., Ritcher, M. J., & Lacy, J. H. 2008, ApJ, 673, 445
Fong, D., Meixner, M., & Shah, R. Y. 2003, ApJ, 582, L39
Guélin, M., Lucas, R., & Cernicharo, J. 1993, A&A, 280, 19
Guélin, M., Lucas, R., & Neri, R. 1997, in IAU Symp. 170, CO: Twenty-Five Years of Millimeter Wave Spectroscopy, ed. W. B. Latter, et al. (Dordrecht: Kluwer), 359
Hasegawa, T. I., et al. 2006, ApJ, 637, 791
Justtanont, K., Skinner, C. J., & Tielens, A. G. G. M. 1994, ApJ, 435, 852
Keady, J. J. & Ridgway, S. T. 1993, ApJ, 406, 199
Leão, I. C., et al. 2006, A&A, 455, 187
Maercker, M., et al. 2008, A&A, 479, 779
Mamon, G. A., Glassgold, A. E., & Huggins, P. J. 1988, ApJ, 328, 797
Melnick, G. J., et al. 2001, Nature, 412, 160
Menten, K. M. & Alcolea, J. 1995, ApJ, 448, 416
Milam, S. N., Woolf, N. J., & Ziurys, L. M. 2009, ApJ, 690, 837
Schöier, F. L., Olofsson, H., & Lundgren, A. A. 2006, A&A, 454, 247
Schöier, F. L., Bast, J., Olofsson, H., & Lindqvist, M. 2007, A&A, 473, 871
Tejero, J., & Cernicharo, J. 1991, IGN, Madrid
Tenenbaum, E. D., Woolf, N. J., & Ziurys, L. M. 2007, ApJ, 666, L29

Teyssier, D., et al. 2006, A&A, 450, 167
Tsuji, T. 1973, A&A, 23, 411
Tsuji, T., Ohnaka, K., Aoki, W., & Yamamura, I. 1997, A&A, 320, L1
van Dishoeck, E. F., Jonkheid, B., & van Hemert, M. C. 2006, Faraday Discuss., 133, 231
Weigelt, G., et al. 1998, A&A, 333, L51
Willacy, K. & Millar, T. J. 1997, A&A, 324, 237
Willacy, K. 2004, ApJ, 600, L87
Woodall, J., Agúndez, M., Markwick-Kemper, A. J., & Millar, T. J. 2007, A&A, 466, 1197
Ziurys, L. M., Milam, S. N., Apponi, A. J., & Woolf, N. J. 2007, Nature, 447, 1094